# Anti-Phospho-eIF4E (Ser209) Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P010154

## Product Information

| | |
|---|---|
| Application | WB, IHC-P/IF (Tissue-P), ICC/IF (Cell), IP, ELISA |
| Reactivity | Human, Mouse, Rat |
| Dilution | WB 1:500~1:1,000; IHC-P 1:50~1:100; IF 1:50~1:200; IP 1:20 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Label | Unconjugated |
| Immunogen | A synthesized peptide derived from human Phospho-eIF4E (S209) |
| Format | Buffer System: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification: Affinity Chromatography |
| Storage | Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage. |
| Precautions | Anti-Phospho-eIF4E (Ser209) antibody is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| Synonyms | EIF4E, EIF4EL1, EIF4F, Eukaryotic translation initiation factor 4E, eIF-4E, eIF4E, eIF-4F 25 kDa subunit, mRNA cap-binding protein. |
| Calculated MW | Calculated MW: 25 kDa; Observed MW: 25 kDa |
| Uniprot ID | P06730 |
| Gene ID | 1977 |
| Background | eIF4E, a protein modulates translation of maternal mRNAs in early embryos before the onset of zygotic transcription. eIF4E also influences the overall rate of translation. eIF4E binds to the 7 methyl GTP cap structure of eukaryotic mRNAs. Phosphorylation of eIF4E on serine 209 regulates the affinity of this protein for the 7 methyl GTP cap and/or RNA. Phosphorylation also enhances the interaction of eIF4E with eIF4G, which form a complex known as eIF4F. eIF4E phosphorylation is correlated with increased translational rate in a number of cell types. |

Western blot analysis of eIF4E (Phospho-S209) in (1) HEK293 lysates; (2) mouse spleen lysates using Phospho-eIF4E (Ser209) antibody.



Immunohistochemistry analysis of paraffin-embedded mouse kidney using Phospho-eIF4E (S209) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.