

Anti-Phospho-IKK alpha (Thr23) Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P013706


Product Information

Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:1,000; IHC-P 1:50~1:100
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Synthetic peptide of human CHUK
Format	Buffer System: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3. Purification: Affinity Purified.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-IKK alpha (Thr23) antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	CHUK, IKKA, TCF16, Inhibitor of nuclear factor kappa-B kinase subunit alpha, I-kappa-B kinase alpha, IKK-A, IKK-alpha, IkbKA, IkappaB kinase, Conserved helix-loop-helix ubiquitous kinase, I-kappa-B kinase 1, IKK1, Nuclear factor NF-kappa-B.
Calculated MW	Calculated MW: 85 kDa; Observed MW: 85 kDa
Uniprot ID	O15111
Gene ID	1147
Background	Plays an essential role in the NF-kappa-B signaling pathway which is activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA damages or other cellular stresses. Activation of IKK depends upon phosphorylation at Ser177 and Ser181 in the activation loop of IKK β (Ser176 and Ser180 in IKK α), which causes conformational changes, resulting in kinase activation.

Validation Images



Western blot analysis of Phospho-IKK alpha (Thr23) in 293 lysates using Phospho-IKK alpha (Thr23) antibody.



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using IKK a(Phospho-Thr23) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Sample with blocking peptide on the right.