

Anti-IKK beta Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P010470


Product Information

Application	WB, IHC-P/IF (Tissue-P), ICC/IF (Cell), ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:1,000; IHC-P 1:50~1:100; IF 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human IKK beta
Format	Buffer System: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification: Affinity Chromatography
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-IKK beta antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	IKKB, IKK2, Inhibitor of nuclear factor kappa-B kinase subunit beta, I-kappa-B-kinase beta, IKK-B, IKK-beta, IkbKB, I-kappa-B kinase 2, IKK2, Nuclear factor NF-kappa-B inhibitor kinase beta, NFKB1KB.
Calculated MW	Calculated MW: 87 kDa; Observed MW: 87 kDa
Uniprot ID	O14920
Gene ID	3551
Background	The NF- κ B/Rel transcription factors are present in the cytosol in an inactive state, complexed with the inhibitory I κ B proteins (1-3). Most agents that activate NF- κ B do so through a common pathway based on phosphorylation-induced, proteasome-mediated degradation of I κ B (3-7). The key regulatory step in this pathway involves activation of a high molecular weight I κ B kinase (IKK) complex whose catalysis is generally carried out by three tightly associated IKK subunits.


Validation Images



Western blot analysis of IKK beta in HeLa lysates using IKK beta antibody.



Immunohistochemistry analysis of paraffin-embedded Human kidney using IKK beta antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunofluorescence analysis of IKK beta in HeLa using IKK beta antibody.