

## Anti-FDFT1 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R013222

### Product Information

Application	WB, IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	20L04L51
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human FDFT1
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-FDFT1 Rabbit mAb [20L04L51] is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information

Synonyms	DGPT, ERG9, Farnesyl diphosphate farnesyltransferase, Farnesyl-diphosphate farnesyltransferase, FDFT_HUMAN, FDFT1, FPP:FPP farnesyltransferase, SQS, Squalene synthase, Squalene synthetase, SS.
Calculated MW	Calculated MW: 48 kDa; Observed MW: 48 kDa
Uniprot ID	P37268
Gene ID	2222
Background	This gene encodes a membrane-associated enzyme located at a branch point in the mevalonate pathway. The encoded protein is the first specific enzyme in cholesterol biosynthesis, catalyzing the dimerization of two molecules of farnesyl diphosphate in a two-step reaction to form squalene. [provided by RefSeq, Jul 2008]
Cellular Location	Endoplasmic reticulum membrane.



Western Blot - Anti-FDFT1 Rabbit mAb [20L04L51]

All lanes: R013222 at 1:1,000 dilution

Lane 1: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 2: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 3: A431 (Human epidermoid teratoma cell line) whole cell lysates

Lane 4: Caco2 (Human colorectal adenocarcinoma epithelial cell) whole cell lysates

Lane 5: SH-SY5Y (Human neuroblastoma epithelial cell) whole cell lysates

Lane 6: 293T (Human embryonic kidney cell) whole cell lysates

Lane 7: SCC-9 (Human tongue squamous carcinoma epithelial cell) whole cell lysates

Lane 8: C2C12 (Mouse myoblasts epithelial cell) whole cell lysates

Lane 9: Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates

Lane 10: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 48 kDa

Observed band size: 48 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-FDFT1 Rabbit mAb [20L04L51]


Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R013222 at 1:100 dilution and  $\alpha$ -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).



Immunohistochemistry - Anti-FDFT1 Rabbit mAb [20L04L51]

Sample: Paraformaldehyde-fixed, paraffin embedded rat stomach tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R013222 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-FDFT1 Rabbit mAb [20L04L51]

Sample: Paraformaldehyde-fixed, paraffin embedded human lung cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R013222 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-FDFT1 Rabbit mAb [20L04L51]

Sample: Paraformaldehyde-fixed, paraffin embedded mouse liver tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R013222 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.