

Anti-Phospho-Nrf2 (Ser40) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R012016

Product Information

Application	ELISA, WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	43K50M15
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Ser40 of human Nrf2
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-Nrf2 (Ser40) Rabbit mAb [43K50M15] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	erythroid derived 2, HEBP1, like 2, NF E2 related factor 2, NF-E2-related factor 2, NF2L2_HUMAN, NFE2 related factor 2, NFE2-related factor 2, Nfe2l2, Nrf 2, NRF2, Nuclear factor (erythroid derived 2) like 2, Nuclear factor, nuclear factor erythroid 2 like 2, Nuclear factor erythroid 2 related factor 2, Nuclear factor erythroid 2-related factor 2, Nuclear factor erythroid derived 2 like 2.
Calculated MW	Calculated MW: 68 kDa; Observed MW: 100 kDa
Uniprot ID	Q16236
Gene ID	4780
Background	This gene encodes a transcription factor which is a member of a small family of basic leucine zipper (bZIP) proteins. The encoded transcription factor regulates genes which contain antioxidant response elements (ARE) in their promoters; many of these genes encode proteins involved in response to injury and inflammation which includes the production of free radicals. Multiple transcript variants encoding different isoforms have been characterized for this gene. [provided by RefSeq, Sep 2015]
Cellular Location	Cytoplasm > cytosol. Nucleus. Cytosolic under unstressed conditions, translocates into the nucleus upon induction by electrophilic agents.
Tissue Location	Widely expressed. Highest expression in adult muscle, kidney, lung, liver and in fetal muscle.



Western Blot - Anti-Phospho-Nrf2 (Ser40) Rabbit mAb [43K50M15]

All lanes: R012016 at 1:2,000 dilution

Lane 1: A431 (Human epidermoid teratoma cell line) whole cell lysates

Lane 2: T24 (Human bladder cancer epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 68 kDa

Observed band size: 100 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-Phospho-Nrf2 (Ser40) Rabbit mAb [43K50M15]

Sample: Paraformaldehyde-fixed, paraffin embedded human cervical cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R012016 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Phospho-Nrf2 (Ser40) Rabbit mAb [43K50M15]

Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R012016 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-Phospho-Nrf2 (Ser40) Rabbit mAb [43K50M15]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R012016 at 1:100 dilution and alpha-tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).