

Anti-ATM Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R010965

Product Information

Application	IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA, WB
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:200; IF 1:100
Host	Rabbit
Clonality	Monoclonal
Clone No.	75K29L94
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human ATM
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-ATM Rabbit mAb [75K29L94] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	ATM, Serine-protein kinase ATM, Ataxia telangiectasia mutated, A-T mutated.
Calculated MW	Calculated MW: 350 kDa; Observed MW: 350 kDa
Uniprot ID	Q13315
Gene ID	472
Background	The protein encoded by this gene belongs to the PI3/PI4-kinase family. This protein is an important cell cycle checkpoint kinase that phosphorylates; thus, it functions as a regulator of a wide variety of downstream proteins, including tumor suppressor proteins p53 and BRCA1, checkpoint kinase CHK2, checkpoint proteins RAD17 and RAD9, and DNA repair protein NBS1. This protein and the closely related kinase ATR are thought to be master controllers of cell cycle checkpoint signaling pathways that are required for cell response to DNA damage and for genome stability.



Western Blot - Anti-ATM Rabbit mAb [75K29L94]

All lanes: R012956 at 1:2,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysates

Lane 3: Jurkat (Human T lymphocytic leukemia cell) whole cell lysates

Lane 4: T24 (Human bladder cancer epithelial cell) whole cell lysates

Lane 5: HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 350 kDa

Observed band size: 350 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-ATM Rabbit mAb [75K29L94]


Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R010965 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (CY3) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).



Immunohistochemistry - Anti-ATM Rabbit mAb [75K29L94]

Sample: Paraformaldehyde-fixed, paraffin embedded human colorectal carcinoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R010965 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-ATM Rabbit mAb [75K29L94]

Sample: Paraformaldehyde-fixed, paraffin embedded human lung cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R010965 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.