

Anti-CDK4 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R012806

Product Information

Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Mouse
Dilution	WB 1:1,000~1:2,000; IHC-P 1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	66L24K34
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human Cdk4
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-CDK4 antibody [66L24K34] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	CMM3, PSK-J3.
Calculated MW	Calculated MW: 33 kDa; Observed MW: 33 kDa
Uniprot ID	P11802
Gene ID	1019
Background	Cyclin-dependent kinase activity is regulated by T-loop phosphorylation (Thr172 in the case of CDK4), by the abundance of their cyclin partners, and by association with CDK inhibitors of the Cip/Kip or INK family of proteins. Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals.



Western Blot - Anti-CDK4 Rabbit mAb [66L24K34]

All lanes: R012806 at 1:1,000 dilution

Lane 1: 293T (human embryonic kidney cell) whole cell lysates

Lane 2: HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 3: HepG2 (human hepatocellular carcinoma epithelial cell) whole cell lysates

Lane 4: SCC-9 (human tongue squamous carcinoma epithelial cell) whole cell lysates

Lane 5: Balb/c mouse liver whole tissue lysates

Lane 6: Balb/c mouse kidney whole tissue lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 33 kDa

Observed band size: 33 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-CDK4 Rabbit mAb [66L24K34]

Sample: Paraformaldehyde-fixed, paraffin embedded rat ovary tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R012806 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-CDK4 Rabbit mAb [66L24K34]

Sample: Paraformaldehyde-fixed, paraffin embedded rat skin tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R012806 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-CDK4 Rabbit mAb [66L24K34]

Sample: Paraformaldehyde-fixed, paraffin embedded human cervical cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R012806 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-CDK4 Rabbit mAb [66L24K34]

Sample: Paraformaldehyde-fixed, paraffin embedded human colorectal carcinoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R012806 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-CDK4 Rabbit mAb [66L24K34]

Sample: Paraformaldehyde-fixed, paraffin embedded mouse subcutaneous tumor tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R012806 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.