

Anti-Phospho-c-Myc (Ser62) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R010019

Product Information

Application	ELISA, IHC-P/IF (Tissue-P), IF (Cell)/ICC, WB
Reactivity	Rat, Human
Dilution	WB 1:4,000; IHC-P 1:200; IF 1:100
Host	Rabbit
Clonality	Monoclonal
Clone No.	62M57L73
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Ser62 of human c-Myc
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-c-Myc (Ser62) Rabbit mAb [62M57L73] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	MYC, BHLHE39, Myc proto-oncogene protein, Class E basic helix-loop-helix protein 39, bHLHe39, Proto-oncogene c-Myc, Transcription factor p64.
Calculated MW	Calculated MW: 48 kDa; Observed MW: 57 kDa
Uniprot ID	P01106
Gene ID	4609
Background	Myc a proto-oncogenic transcription factor that plays a role in cell proliferation, apoptosis and in the development of human tumors.. Seems to activate the transcription of growth-related genes.



Western Blot - Anti-Phospho-c-Myc (Ser62) Rabbit mAb [62M57L73]

All lanes: R010019 at 1:4,000 dilution

Lane 1: A431 (human epidermoid carcinoma cell line) whole cell lysates

Lane 2: Jurkat (human T lymphocytic leukemia cell) whole cell lysates

Lane 3: HCT116 (human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: SW620 (human colorectal carcinoma epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 48 kDa

Observed band size: 57 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-Phospho-c-Myc (Ser62) Rabbit mAb [62M57L73]

Sample: Paraformaldehyde-fixed, paraffin embedded rat testis tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R010019 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Phospho-c-Myc (Ser62) Rabbit mAb [62M57L73]

Sample: Paraformaldehyde-fixed, paraffin embedded human colorectal carcinoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R010019 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-Phospho-c-Myc (Ser62) Rabbit mAb [62M57L73]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R010019 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and

Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).