

Anti-ELK1 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R010378

Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	26M38M98
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human ELK1
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-ELK1 Rabbit mAb [26M38M98] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	ETS domain-containing protein Elk-1; ELK1.
Calculated MW	Calculated MW: 45 kDa; Observed MW: 62 kDa
Uniprot ID	P19419
Gene ID	2002
Background	This gene is a member of the Ets family of transcription factors and of the ternary complex factor (TCF) subfamily. Proteins of the TCF subfamily form a ternary complex by binding to the the serum response factor and the serum response element in the promoter of the c-fos proto-oncogene. The protein encoded by this gene is a nuclear target for the ras-raf-MAPK signaling cascade. This gene produces multiple isoforms by using alternative translational start codons and by alternative splicing. Related pseudogenes have been identified on chromosomes 7 and 14. [provided by RefSeq, Mar 2012]
Cellular Location	Nucleus
Tissue Location	Lung and testis.



Western Blot - Anti-ELK1 Rabbit mAb [26M38M98]

All lanes: R010378 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: K562 (Human chronic myeloid leukemia cell) whole cell lysates

Lane 3: U87 (Human malignant glioblastoma epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 45 kDa

Observed band size: 62 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-ELK1 Rabbit mAb [26M38M98]


Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R010378 at 1:100 dilution and alpha-tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).



Immunohistochemistry - Anti-ELK1 Rabbit mAb [26M38M98]

Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R010378 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.