

Anti-FAK Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R012430

Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Mouse, Rat, Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:200; IF 1:100
Host	Rabbit
Clonality	Monoclonal
Clone No.	56K44M73
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human FAK
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-FAK antibody [56K44M73] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	PTK2, FAK, FAK1, Focal adhesion kinase 1, FADK 1, Focal adhesion kinase-related nonkinase, FRNK, Protein phosphatase 1 regulatory subunit 71, PPP1R71, Protein-tyrosine kinase 2, p125FAK, pp125FAK.
Calculated MW	Calculated MW: 119 kDa; Observed MW: 120-130 kDa
Uniprot ID	Q05397
Gene ID	5747
Background	This gene encodes a cytoplasmic protein tyrosine kinase which is found concentrated in the focal adhesions that form between cells growing in the presence of extracellular matrix constituents. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Activation of this gene may be an important early step in cell growth and intracellular signal transduction pathways triggered in response to certain neural peptides or to cell interactions with the extracellular matrix.



Western Blot - Anti-FAK Rabbit mAb [56K44M73]

All lanes: R012430 at 1:1,000 dilution

Lane 1: HCT116 (human colorectal carcinoma epithelial cell) whole cell lysates

Lane 2: MCF7 (human breast adenocarcinoma epithelial cell) whole cell lysates

Lane 3: T24 (human bladder cancer epithelial cell) whole cell lysates

Lane 4: Rat brainwhole tissue lysates

Lane 5: Rat ovarywhole tissue lysates

Lane 6: Balb/c mouse brain whole tissue lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 119 kDa

Observed band size: 120-130 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-FAK Rabbit mAb [56K44M73]

Sample: Paraformaldehyde-fixed, paraffin embedded human colorectal carcinoma tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R012430 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-FAK Rabbit mAb [56K44M73]

Sample: Paraformaldehyde-fixed, paraffin embedded mouse kidney tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R012430 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-FAK Rabbit mAb [56K44M73]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R012430 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).