

## Anti-Hsp90 alpha/beta Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R010492

### Product Information

Application	IF (Cell)/ICC, ELISA, WB, IHC-P/IF (Tissue-P)
Reactivity	Mouse, Rat, Human
Dilution	WB 1:1,000~1:6,000; IHC-P 1:200; IF 1:100
Host	Rabbit
Clonality	Monoclonal
Clone No.	26M41K41
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human Hsp90 beta
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Hsp90 alpha/beta Rabbit mAb [26M41K41] is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information

Synonyms	HSP90AA1, HSP90A, HSPC1, HSPCA, Heat shock protein HSP 90-alpha, Heat shock 86 kDa, HSP 86, HSP86, Renal carcinoma antigen NY-REN-38, HSP90AB1, HSP90B, HSPC2, HSPCB, Heat shock protein HSP 90-beta, HSP 90, Heat shock 84 kDa, HSP 84, HSP84.
Calculated MW	Calculated MW: 90 kDa; Observed MW: 90 kDa
Uniprot ID	P07900, P08238
Gene ID	3320/3326
Background	Molecular chaperone that promotes the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction. Undergoes a functional cycle that is linked to its ATPase activity. This cycle probably induces conformational changes in the client proteins, thereby causing their activation.



Western Blot - Anti-Hsp90 alpha/beta Rabbit mAb [26M41K41]

All lanes: R010492 at 1:6,000 dilution

Lane 1: U2OS (Human osteosarcoma epithelial cell) whole cell lysates

Lane 2: Rat heart whole cell lysates

Lane 3: Rat pancreas whole cell lysates

Lane 4: Balb/c mouse kidney whole tissue lysates

Lane 5: Balb/c mouse brain whole tissue lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 90 kDa

Observed band size: 90 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-Hsp90 alpha/beta Rabbit mAb [26M41K41]


Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R010492 at 1:100 dilution and  $\alpha$ -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).



Immunohistochemistry - Anti-Hsp90 alpha/beta Rabbit mAb [26M41K41]

Sample: Paraformaldehyde-fixed, paraffin embedded human breast carcinoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R010492 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Hsp90 alpha/beta Rabbit mAb [26M41K41]

Sample: Paraformaldehyde-fixed, paraffin embedded rat testis tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R010492 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Hsp90 alpha/beta Rabbit mAb [26M41K41]

Sample: Paraformaldehyde-fixed, paraffin embedded mouse uterus tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R010492 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.