

## Anti-MEF2A Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R013467

### Product Information

Application	IF (Cell)/ICC, WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:3,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	84L74M32
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human MEF2A
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-MEF2A Rabbit mAb [84L74M32] is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information

Synonyms	ADCAD1, MADS box transcription enhancer factor 2, polypeptide A (myocyte enhancer factor 2A), MEF2, MEF2A, MEF2A_HUMAN, Myocyte enhancer factor 2A, Myocyte-specific enhancer factor 2A, RSRFC4, RSRFC9, Serum response factor like protein 1, Serum response factor-like protein 1.
Calculated MW	Calculated MW: 55 kDa; Observed MW: 55 kDa
Uniprot ID	Q02078
Gene ID	4205
Background	Transcriptional activator which binds specifically to the MEF2 element, 5'-YTA[AT]4TAR-3', found in numerous muscle-specific genes. Also involved in the activation of numerous growth factor- and stress-induced genes. Mediates cellular functions not only in skeletal and cardiac muscle development, but also in neuronal differentiation and survival. Plays diverse roles in the control of cell growth, survival and apoptosis via p38 MAPK signaling in muscle-specific and/or growth factor-related transcription. In cerebellar granule neurons, phosphorylated and sumoylated MEF2A represses transcription of NUR77 promoting synaptic differentiation. Associates with chromatin to the ZNF16 promoter.
Cellular Location	Nucleus.
Tissue Location	Isoform MEF2 and isoform MEFA are expressed only in skeletal and cardiac muscle and in the brain. Isoform RSRFC4 and isoform RSRFC9 are expressed in all tissues examined.



Western Blot - Anti-MEF2A Rabbit mAb [84L74M32]

All lanes: R013467 at 1:3,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: T24 (Human bladder cancer epithelial cell) whole cell lysates

Lane 3: Rat heart whole tissue lysates

Lane 4: Mouse muscle whole tissue lysates


Lysates/proteins at 10  $\mu$ g per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 55 kDa

Observed band size: 55 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-MEF2A Rabbit mAb [84L74M32]

Sample: Paraformaldehyde-fixed, paraffin embedded rat liver tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R013467 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-MEF2A Rabbit mAb [84L74M32]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R013467 at 1:100 dilution and  $\alpha$ -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).