

Anti-Phospho-MEK1 (Ser298) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R012258

Product Information

Application	ELISA, WB, IF (Cell)/ICC
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	36M99L32
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Ser298 of human MEK1
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-MEK1 (Ser298) Rabbit mAb [36M99L32] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Dual specificity mitogen activated protein kinase kinase 1, Dual specificity mitogen-activated protein kinase kinase 1, ERK activator kinase 1, MAP kinase kinase 1, MAP kinase/Erk kinase 1, MAP2K1, MAPK/ERK kinase 1, MAPKK 1, MAPKK1, MEK 1, Mek1, MEKK1, Mitogen activated protein kinase kinase 1, MKK 1, MKK1, MP2K1_HUMAN, PRKMK1, Protein kinase mitogen activated kinase 1 (MAP kinase kinase 1), Protein kinase mitogen activated, kinase 1, protein kinase mitogen-activated kinase 1.
Calculated MW	Calculated MW: 43 kDa; Observed MW: 43 kDa
Uniprot ID	Q02750
Gene ID	5604
Background	The protein encoded by this gene is a member of the dual specificity protein kinase family, which acts as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein kinase lies upstream of MAP kinases and stimulates the enzymatic activity of MAP kinases upon wide variety of extra- and intracellular signals. As an essential component of MAP kinase signal transduction pathway, this kinase is involved in many cellular processes such as proliferation, differentiation, transcription regulation and development. [provided by RefSeq, Jul 2008]
Tissue Location	Widely expressed, with extremely low levels in brain.



Western Blot - Anti-Phospho-MEK1 (Ser298) Rabbit mAb [36M99L32]

All lanes: R012258 at 1:1,000 dilution

Lane 1: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 2: SCC-9 (Human tongue squamous carcinoma epithelial cell) whole cell lysates

Lane 3: SW620 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: C2C12 (Mouse myoblasts epithelial cell) whole cell lysates

Lane 5: HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates

Lane 6: Balb/c mouse brain whole tissue lysates


Lysates/proteins at 10 μ g per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 43 kDa

Observed band size: 43 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-Phospho-MEK1 (Ser298) Rabbit mAb [36M99L32]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R012258 at 1:100 dilution and α -tubulin Mouse Monoclonal

Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and

Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).