

Anti-SCD1 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R010079

Product Information

Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	60L70M15
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human SCD1
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-SCD1 Rabbit mAb [60L70M15] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	FADS5; SCD1; SCDOS; SCD; Stearoyl-CoA desaturase; hSCD1; Acyl-CoA desaturase; Delta(9)-desaturase; Fatty acid desaturase; Delta-9 desaturase.
Calculated MW	Calculated MW: 42 kDa; Observed MW: 42 kDa
Uniprot ID	O00767
Gene ID	6319
Background	This gene encodes an enzyme involved in fatty acid biosynthesis, primarily the synthesis of oleic acid. The protein belongs to the fatty acid desaturase family and is an integral membrane protein located in the endoplasmic reticulum. Transcripts of approximately 3.9 and 5.2 kb, differing only by alternative polyadenylation signals, have been detected. A gene encoding a similar enzyme is located on chromosome 4 and a pseudogene of this gene is located on chromosome 17. [provided by RefSeq, Sep 2015]
Cellular Location	Endoplasmic reticulum membrane.Multi-pass membrane protein.
Tissue Location	Detected in fetal liver, lung and brain. Highly expressed in adult adipose tissue, and at lower levels in adult brain and lung.



Western Blot - Anti-SCD1 mAb [60L70M15]

All lanes: R010079 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: A431 (Human epidermoid teratoma cell line) whole cell lysates

Lane 5: U87 (Human malignant glioblastoma epithelial cells) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 42 kDa

Observed band size: 42 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-SCD1 mAb [60L70M15]

Sample: Paraformaldehyde-fixed, paraffin embedded human hepatoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R010079 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-SCD1 mAb [60L70M15]

Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R010079 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.