

Anti-XIAP Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R011715

Product Information

Application	WB, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000
Host	Rabbit
Clonality	Monoclonal
Clone No.	41L25L98
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human XIAP
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-XIAP Rabbit mAb [41L25L98] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Aipa; Api3; Birc4; Miha; Xiap; E3 ubiquitin-protein ligase XIAP; Baculoviral IAP repeat-containing protein 4; IAP homolog A; Inhibitor of apoptosis protein 3; RING-type E3 ubiquitin transferase XIAP; X-linked inhibitor of apoptosis protein; IAP-3; mIAP-3; mIAP3; X-linked IAP.
Calculated MW	Calculated MW: 57 kDa; Observed MW: 57 kDa
Uniprot ID	Q60989
Gene ID	11798
Background	This gene encodes a protein that belongs to a family of apoptotic suppressor proteins. Members of this family share a conserved motif termed, baculovirus IAP repeat, which is necessary for their anti-apoptotic function. This protein functions through binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2 and inhibits apoptosis induced by menadione, a potent inducer of free radicals, and interleukin 1-beta converting enzyme. This protein also inhibits at least two members of the caspase family of cell-death proteases, caspase-3 and caspase-7. Mutations in this gene are the cause of X-linked lymphoproliferative syndrome. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 2 and 11.[provided by RefSeq, Feb 2011]
Cellular Location	Cytoplasm.Nucleus.TLE3 promotes its nuclear localization.
Tissue Location	Expressed in small intestine crypts (at protein level) (PubMed:30389919). Expressed in bulge hair follicle stem cells, sebaceous glands and dermal papillae (at protein level) (PubMed:23788729).



Western Blot - Anti-XIAP Rabbit mAb [41L25L98]

All lanes: R011715 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: 293T (Human embryonic kidney cell) whole cell lysates

Lane 5: K562 (Human chronic myeloid leukemia cell) whole cell lysates

Lane 6: SH-SY5Y (Human neuroblastoma epithelial cell) whole cell lysates

Lane 7: C2C12 (Mouse myoblasts epithelial cell) whole cell lysates

Lane 8: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates

Lane 9: Rat brain whole tissue lysates

Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 57 kDa

Observed band size: 57 kDa

Developed using the ECL technique (Cat. No. SQ201).