

Anti-APC Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R011345

Product Information

Application	WB, ELISA
Reactivity	Mouse, Rat, Human
Dilution	WB 1:1,000~1:2,000
Host	Rabbit
Clonality	Monoclonal
Clone No.	53L68M24
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic peptide of human APC
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-APC antibody [53L68M24] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Adenomatous Polyposis Coli, Adenomatous polyposis coli protein, Apc, APC_HUMAN, CC1, Deleted in polyposis 2.5, DP2, DP2.5, DP3, FPC, GS, Protein APC.
Calculated MW	Calculated MW: 312 kDa; Observed MW: 160 kDa
Uniprot ID	P25054
Gene ID	324
Background	Tumor suppressor. Promotes rapid degradation of CTNNB1 and participates in Wnt signaling as a negative regulator. APC activity is correlated with its phosphorylation state. Activates the GEF activity of SPATA13 and ARHGEF4. Plays a role in hepatocyte growth factor (HGF)-induced cell migration. Required for MMP9 up-regulation via the JNK signaling pathway in colorectal tumor cells. Acts as a mediator of ERBB2-dependent stabilization of microtubules at the cell cortex.
Cellular Location	Cell junction > adherens junction. Cytoplasm > cytoskeleton. Cell projection > lamellipodium. Cell projection > ruffle membrane. Cytoplasm. Cell membrane. Associated with the microtubule network at the growing distal tip of microtubules. Accumulates in the lamellipodium and ruffle membrane in response to hepatocyte growth factor (HGF) treatment. The MEMO1-RHOA-DIAPH1 signaling pathway controls localization of the phosphorylated form to the cell membrane.
Tissue Location	Expressed in a variety of tissues.



Western Blot - Anti-APC Rabbit mAb [53L68M24]

All lanes: R011345 at 1:1,000 dilution

Lane 1: C2C12 (Mouse myoblasts epithelial cell) whole cell lysates

Lane 2: U2OS (Human osteosarcoma epithelial cell) whole cell lysates

Lane 3: T24 (Human bladder cancer epithelial cell) whole cell lysates

Lane 4: Rat heart whole tissue lysates

Lane 5: Rat brain whole tissue lysates

Lane 6: Rat spleen whole tissue lysates

Lane 7: Mouse small intestine whole tissue lysates

Lysates/proteins at 10 μ g per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 312 kDa

Observed band size: 160 kDa

Developed using the ECL technique (Cat. No. SQ201).