

Anti-Daxx Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R011698

Product Information

Application	ELISA, WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	23M45K41
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Daxx
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Daxx Rabbit mAb [23M45K41] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	BING 2,BING2,CENP-C binding protein,DAP 6,DAP6,Daxx,DAXX_HUMAN,Death associated protein 6,Death domain associated protein 6,Death domain associated protein,Death domain-associated protein 6,EAP 1,EAP1,ETS1 associated protein 1,ETS1-associated protein 1,Fas binding protein,Fas death domain associated protein,Fas death domain-associated protein,hDaxx,MGC126245,MGC126246.
Calculated MW	Calculated MW: 81 kDa; Observed MW: 110 kDa
Uniprot ID	Q9UER7
Gene ID	1616
Background	This gene encodes a multifunctional protein that resides in multiple locations in the nucleus and in the cytoplasm. It interacts with a wide variety of proteins, such as apoptosis antigen Fas, centromere protein C, and transcription factor erythroblastosis virus E26 oncogene homolog 1. In the nucleus, the encoded protein functions as a potent transcription repressor that binds to sumoylated transcription factors. Its repression can be relieved by the sequestration of this protein into promyelocytic leukemia nuclear bodies or nucleoli. This protein also associates with centromeres in G2 phase. In the cytoplasm, the encoded protein may function to regulate apoptosis. The subcellular localization and function of this protein are modulated by post-translational modifications, including sumoylation, phosphorylation and polyubiquitination. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2008]
Cellular Location	Nucleus. Diffuse nuclear distribution pattern and no comparable dot-like accumulation of isoform 1 and Cytoplasm. Nucleus, nucleoplasm. Nucleus, PML body. Nucleus, nucleolus. Chromosome, centromere. Dispersed throughout the nucleoplasm, in PMI/POD/ND10 nuclear bodies. and in nucleoli (Probable). Colocalizes with histone H3.3. ATRX. HIRA and ASF1A at PMI-



Western Blot - Anti-Daxx Rabbit mAb [23M45K41]

All lanes: R011698 at 1:2,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: A431 (Human epidermoid teratoma cell line) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 81 kDa

Observed band size: 110 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-Daxx Rabbit mAb [23M45K41]

Sample: Paraformaldehyde-fixed, paraffin embedded human colonic cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R011698 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Daxx Rabbit mAb [23M45K41]

Sample: Paraformaldehyde-fixed, paraffin embedded human gastric cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R011698 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-Daxx Rabbit mAb [23M45K41]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R011698 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and

Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).