

## Anti-MK3 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R012293

### Product Information

Application	WB, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:5,000
Host	Rabbit
Clonality	Monoclonal
Clone No.	12L57K12
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human MK-3
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-MK3 Rabbit mAb [12L57K12] is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information

Synonyms	MAP kinase-activated protein kinase 3; MAPK-activated protein kinase 3; MAPKAP kinase 3; MAPKAP-K3; MAPKAPK-3; MK-3; Chromosome 3p kinase; 3pK; MAPKAPK3.
Calculated MW	Calculated MW: 43 kDa; Observed MW: 43 kDa
Uniprot ID	Q16644
Gene ID	7867
Background	This gene encodes a member of the Ser/Thr protein kinase family. This kinase functions as a mitogen-activated protein kinase (MAP kinase)- activated protein kinase. MAP kinases are also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This kinase was shown to be activated by growth inducers and stress stimulation of cells. In vitro studies demonstrated that ERK, p38 MAP kinase and Jun N-terminal kinase were all able to phosphorylate and activate this kinase, which suggested the role of this kinase as an integrative element of signaling in both mitogen and stress responses. This kinase was reported to interact with, phosphorylate and repress the activity of E47, which is a basic helix-loop-helix transcription factor known to be involved in the regulation of tissue-specific gene expression and cell differentiation. Alternate splicing results in multiple transcript variants that encode the same protein. [provided by RefSeq, Sep 2011]
Cellular Location	Nucleus.Cytoplasm.Predominantly located in the nucleus, when activated it translocates to the cytoplasm.
Tissue Location	Widely expressed, with a higher expression level observed in heart and skeletal muscle. No expression in brain. Expressed in the retinal pigment epithelium (PubMed:26744326).



Western Blot - Anti-MK3 Rabbit mAb [12L57K12]

All lanes: R012293 at 1:5,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: A431 (Human epidermoid teratoma cell line) whole cell lysates

Lane 5: K562 (Human chronic myeloid leukemia cell) whole cell lysates

Lane 6: U87 (Human malignant glioblastoma epithelial cell) whole cell lysates

Lane 7: C2C12 (Mouse myoblasts epithelial cell) whole cell lysates

Lane 8: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates

Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 43 kDa

Observed band size: 43 kDa

Developed using the ECL technique (Cat. No. SQ201).