

Anti-TRIM24 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R013440

Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Mouse, Human, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	44M94M33
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human TRIM24
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-TRIM24 Rabbit mAb [44M94M33] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	E3 ubiquitin protein ligase TRIM24, E3 ubiquitin-protein ligase Trim24, hTIF1, PTC6, RING finger protein 82, RNF82, TF1A, TIF1 alpha, TIF1, TIF1-alpha, TIF1A, TIF1A_HUMAN, TIF1ALPHA, Transcription intermediary factor 1-alpha, Transcriptional intermediary factor 1 alpha, Transcriptional intermediary factor 1, Trim24, Tripartite motif containing 24, Tripartite motif-containing protein 24.
Calculated MW	Calculated MW: 117 kDa; Observed MW: 130-140 kDa
Uniprot ID	O15164
Gene ID	8805
Background	The protein encoded by this gene mediates transcriptional control by interaction with the activation function 2 (AF2) region of several nuclear receptors, including the estrogen, retinoic acid, and vitamin D3 receptors. The protein localizes to nuclear bodies and is thought to associate with chromatin and heterochromatin-associated factors. The protein is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains - a RING, a B-box type 1 and a B-box type 2 - and a coiled-coil region. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008]
Cellular Location	Nucleus. Cytoplasm. Colocalizes with sites of active transcription. Detected both in nucleus and cytoplasm in some breast cancer samples. Predominantly nuclear.



Western Blot - Anti-TRIM24 Rabbit mAb [44M94M33]

All lanes: R013440 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: A431 (Human epidermoid teratoma cell line) whole cell lysates

Lane 5: T24 (Human bladder cancer epithelial cell) whole cell lysates

Lane 6: Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates

Lane 7: U2OS (Human osteosarcoma epithelial cell) whole cell lysates

Lane 8: Rat spleen whole tissue lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 117 kDa

Observed band size: 130-140 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-TRIM24 Rabbit mAb [44M94M33]

Sample: Paraformaldehyde-fixed, paraffin embedded human lung cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R013440 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-TRIM24 Rabbit mAb [44M94M33]

Sample: Paraformaldehyde-fixed, paraffin embedded mouse ovary tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R013440 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-TRIM24 Rabbit mAb [44M94M33]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R013440 at 1:100 dilution and α-tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).