

## Anti-Phospho-PKC (Ser660) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R012953

### Product Information


Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:5,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	68K33K11
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Ser660 of human PKC
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-PKC (Ser660) Rabbit mAb [68K33K11] is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information


Synonyms	KPCA_HUMAN; PKC alpha; PKC beta; PKC delta; PKC epsilon; PKC gamma; PKC zeta; PKC-A; PKC-alpha; PKC2; PKCA; PKCB; PKCD; PKCE; PKCG; PRKCA; PRKCB; PRKCB1; PRKCB2; PRKCD; PRKCE; PRKCG; PRKCZ; Protein kinase C alpha; Protein kinase C alpha type; Protein kinase C; Protein kinase C beta; Protein kinase C delta; Protein kinase C epsilon; Protein kinase C gamma; Protein kinase C zeta.
Calculated MW	Calculated MW: 78 kDa; Observed MW: 78-85 kDa
Uniprot ID	P17252, P05771, P05771-2, Q05655, Q02156, P24723, Q04759
Gene ID	5578, 5579, 5580, 5581, 5583, 5588
Background	Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This kinase has been reported to play roles in many different cellular processes, such as cell adhesion, cell transformation, cell cycle checkpoint, and cell volume control. Knockout studies in mice suggest that this kinase may be a fundamental regulator of cardiac contractility and Ca(2+) handling in myocytes. [provided by RefSeq, Jul 2008].
Cellular Location	Cytoplasm. Cell membrane. Mitochondrion membrane. Nucleus.



Immunohistochemistry - Anti-Phospho-PKC (Ser660) Rabbit mAb [68K33K11]  
 Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue  
 Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.  
 Primary antibody: R012953 at 1:200 dilution  
 Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,001 dilution  
 DAB was used as the chromogen.  
 Counter stained with hematoxylin.  
 Positive/negative staining were presented.  
 Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Phospho-PKC (Ser660) Rabbit mAb [68K33K11]  
 Sample: Paraformaldehyde-fixed, paraffin embedded mouse ovary tissue  
 Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.  
 Primary antibody: R012953 at 1:200 dilution  
 Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,001 dilution  
 DAB was used as the chromogen.  
 Counter stained with hematoxylin.  
 Positive/negative staining were presented.  
 Only the secondary antibody was used as the negative control.



Western Blot - Anti-Phospho-PKC (Ser660) Rabbit mAb [68K33K11]  
 All lanes: R012953 at 1:5,000 dilution  
 Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates  
 Lane 2: Huh1 (Human hepatocarcinoma epithelial cell) whole cell lysates  
 Lane 3: T24 (Human bladder cancer epithelial cell) whole cell lysates  
 Lane 4: U87 (Human malignant glioblastoma epithelial cells) whole cell lysates  
 Lane 5: Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates  
 Lane 6: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates  
 Lysates/proteins at 10 µg per lane.  
 Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution  
 Predicted band size: 78 kDa  
 Observed band size: 78-85 kDa  
 Developed using the ECL technique (Cat. No. SQ201).