

Anti-Cyclin E1 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R013802

Product Information

Application	IF (Cell)/ICC, ELISA, WB, IHC-P/IF (Tissue-P)
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	56E11C04
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Cyclin E1
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Cyclin E1 Rabbit mAb [56E11C04] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	CCNE, Ccne1, CCNE1_HUMAN, cyclin E variant ex5del, cyclin E variant ex7del, Cyclin E1, Cyclin Es, Cyclin Et, CyclinE, G1/S specific cyclin E, G1/S-specific cyclin-E1.
Calculated MW	Calculated MW: 47 kDa; Observed MW: 47 kDa
Uniprot ID	P24864
Gene ID	898
Background	<p>The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB. [provided by RefSeq, Apr 2016]</p>
Cellular Location	Nucleus.
Tissue Location	Highly expressed in testis and placenta. Low levels in bronchial epithelial cells.



Western Blot - Anti-Cyclin E1 Rabbit mAb [18K76M38]

All lanes: R011802 at 1:1,000 dilution

Lane 1: A549 (human non-small cell lung cancer epithelial cell) whole cell lysates

Lane 2: Jurkat (human T lymphocytic leukemia cell) whole cell lysates

Lane 3: HCT116 (human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: HepG2 (human hepatocellular carcinoma epithelial cell) whole cell lysates

Lane 5: U2OS (human osteosarcoma epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 47 kDa

Observed band size: 47 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-Cyclin E1 Rabbit mAb [18K76M38]

Sample: Paraformaldehyde-fixed, paraffin embedded rat testis tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R013802 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Cyclin E1 Rabbit mAb [18K76M38]

Sample: Paraformaldehyde-fixed, paraffin embedded human colorectal carcinoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R013802 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Cyclin E1 Rabbit mAb [18K76M38]

Sample: Paraformaldehyde-fixed, paraffin embedded rat ovary tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R013802 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-Cyclin E1 Rabbit mAb [56E11C04]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R010294 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).