

Anti-STAT1 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R013811

Product Information

Application	IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA, WB
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	26C92E46
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human STAT1
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-STAT1 Rabbit mAb [26C92E46] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Signal transducer and activator of transcription 1 91kD, CANDF7, DKFZp686B04100, IMD31A, IMD31B, IMD31C, ISGF 3, ISGF-3, OTTHUMP00000163552, OTTHUMP00000165046, OTTHUMP00000165047, OTTHUMP00000205845, Signal transducer and activator of transcription 1 91kDa, Signal transducer and activator of transcription 1, Signal transducer and activator of transcription 1, 91kD, Signal transducer and activator of transcription 1-alpha/beta, STAT 1, Stat1, STAT1_HUMAN, STAT91, Transcription factor ISGF 3 components p91 p84, Transcription factor ISGF-3 components p91/p84, Transcription factor ISGF3 components p91/p84, XStat1.
Calculated MW	Calculated MW: 87 kDa; Observed MW: 87 kDa
Uniprot ID	P42224
Gene ID	6772
Background	The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. Two alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008]
Cellular Location	Cytoplasm, Nucleus. Translocated into the nucleus in response to IFN-gamma-induced tyrosine phosphorylation and dimerization.



Western Blot - Anti-STAT1 Rabbit mAb [26C92E46]

All lanes: R013811 at 1:1,000 dilution

Lane 1: Jurkat (human T lymphocytic leukemia cell) whole cell lysates

Lane 2: HCT116 (human colorectal carcinoma epithelial cell) whole cell lysates

Lane 3: RAW264.7 (mouse mononuclear macrophage leukemia epithelial cell)

Lane 4: HepG2 (human hepatocellular carcinoma epithelial cell) whole cell lysates

Lane 5: SW620 (human colorectal carcinoma epithelial cell) whole cell lysates

Lane 6: Rat kidney whole tissue lysates

Lane 7: Balb/c mouse brain whole tissue lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 87 kDa

Observed band size: 87 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-STAT1 Rabbit mAb [26C92E46]


Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R013811 at 1:50 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:50 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).



Immunohistochemistry - Anti-STAT1 Rabbit mAb [26C92E46]

Sample: Paraformaldehyde-fixed, paraffin embedded mouse brain tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R013811 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-STAT1 Rabbit mAb [26C92E46]

Sample: A431 cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R013811 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).