

Anti-Phospho-CREB (Ser133) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R013837

Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:200; IF 1:100
Host	Rabbit
Clonality	Monoclonal
Clone No.	83G12F48
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Ser133 of human CREB
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-CREB (Ser133) Rabbit mAb [83G12F48] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	CREB1, Cyclic AMP-responsive element-binding protein 1, CREB-1, cAMP-responsive element-binding protein 1, Active transcription factor CREB, cAMP response element binding protein 1, cAMP response element binding protein antibody, MGC9284, Transactivator protein, OTTHUMP00000163864, OTTHUMP00000163865, OTTHUMP00000206660, OTTHUMP00000206662, OTTHUMP00000206667.
Calculated MW	Calculated MW: 46 kDa; Observed MW: 46 kDa
Uniprot ID	P16220
Gene ID	1385
Background	This gene encodes a transcription factor that is a member of the leucine zipper family of DNA binding proteins. This protein binds as a homodimer to the cAMP-responsive element, an octameric palindrome.
Cellular Location	Nucleus.



Western Blot - Anti-Phospho-CREB (Ser133) Rabbit mAb [83G12F48]

All lanes: R013837 at 1:1,000 dilution

Lane 1: Jurkat (Human T lymphocytic leukemia cell) whole cell lysates

Lane 2: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 3: U2OS (Human osteosarcoma epithelial cell) whole cell lysates

Lane 4: SW620 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 5: HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 46 kDa

Observed band size: 46 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-Phospho-CREB (Ser133) Rabbit mAb [83G12F48]

Sample: Paraformaldehyde-fixed, paraffin embedded human breast carcinoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R013837 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-Phospho-CREB (Ser133) Rabbit mAb [83G12F48]


Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R013837 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).



Immunohistochemistry - Anti-Phospho-CREB (Ser133) Rabbit mAb [83G12F48]

Sample: Paraformaldehyde-fixed, paraffin embedded rat colon tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R013837 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Phospho-CREB (Ser133) Rabbit mAb [83G12F48]
Sample: Paraformaldehyde-fixed, paraffin embedded mouse brain tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R013837 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.