

Anti-Acetyl-p53 (Lys382) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R013972

Product Information

Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	89D56L95
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human p53 (acetyl K382)
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Acetyl-p53 (Lys382) Rabbit mAb [89D56L95] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Antigen NY-CO-13, BCC7, Cellular tumor antigen p53, FLJ92943, LFS1, Mutant tumor protein 53, p53, p53 tumor suppressor, P53_HUMAN, Phosphoprotein p53, Tp53, Transformation related protein 53, TRP53, tumor antigen p55, Tumor protein 53, Tumor protein p53, Tumor suppressor p53.
Calculated MW	Calculated MW: 53 kDa; Observed MW: 53 kDa
Uniprot ID	P04637
Gene ID	7157
Background	Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression.
Cellular Location	Cytoplasm; Cytoplasm. Nucleus. Nucleus > PML body. Endoplasmic reticulum. Interaction with BANP promotes nuclear localization. Recruited into PML bodies together with CHEK2; Nucleus. Cytoplasm. Localized in both nucleus and cytoplasm in most cells. In some cells, forms foci in the nucleus that are different from nucleoli; Nucleus. Cytoplasm. Localized in the nucleus in most cells but found in the cytoplasm in some cells; Nucleus. Cytoplasm. Localized mainly in the nucleus with minor staining in the cytoplasm; Nucleus. Cytoplasm. Predominantly nuclear but localizes to the cytoplasm when expressed with isoform 4 and Nucleus. Cytoplasm. Predominantly nuclear but translocates to the cytoplasm following cell stress.



Western Blot - Anti-Acetyl-p53 (Lys382) Rabbit mAb [89D56L95]

All lanes: R013972 at 1:1,000 dilution

Lane 1: A431 (Human epidermoid teratoma cell line) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 53 kDa

Observed band size: 53 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-Acetyl-p53 (Lys382) Rabbit mAb [89D56L95]

Sample: Paraformaldehyde-fixed, paraffin embedded human colorectal carcinoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R013972 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Acetyl-p53 (Lys382) Rabbit mAb [89D56L95]

Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R013972 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Acetyl-p53 (Lys382) Rabbit mAb [89D56L95]

Sample: Paraformaldehyde-fixed, paraffin embedded mouse skin tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R013972 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.