

Anti-IQGAP2 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R014015

Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Mouse, Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	49T11N85
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human IQGAP2
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-IQGAP2 Rabbit mAb [49T11N85] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	IQ motif containing GTPase activating protein 2, IQGA2_HUMAN, Iqgap2, Ras GTPase activating-like protein, Ras GTPase-activating-like protein IQGAP2.
Calculated MW	Calculated MW: 181 kDa; Observed MW: 181 kDa
Uniprot ID	Q13576
Gene ID	10788
Background	Binds to activated CDC42 and RAC1 but does not seem to stimulate their GTPase activity. Associates with calmodulin.
Tissue Location	Isoform 2 expression is enhanced in testis.



Western Blot - Anti-IQGAP2 Rabbit mAb [49T11N85]

All lanes: R014015 at 1:1,000 dilution

Lane 1: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 181 kDa

Observed band size: 181 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-IQGAP2 Rabbit mAb [49T11N85]

Sample: Paraformaldehyde-fixed, paraffin embedded human hepatoma cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R014015 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-IQGAP2 Rabbit mAb [49T11N85]

Sample: Paraformaldehyde-fixed, paraffin embedded human cervical cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R014015 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-IQGAP2 Rabbit mAb [49T11N85]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R014015 at 1:100 dilution and alpha-tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and

Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).