

Anti-MyD88 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R014035

Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:200; IF 1:100
Host	Rabbit
Clonality	Monoclonal
Clone No.	43Q70G58
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human MyD88
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-MyD88 Rabbit mAb [43Q70G58] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Myeloid differentiation primary response protein MyD88, MYD88.
Calculated MW	Calculated MW: 33 kDa; Observed MW: 33 kDa
Uniprot ID	Q99836
Gene ID	4615
Background	Members of the Toll-like receptor (TLR) family, named for the closely related Toll receptor in <i>Drosophila</i> , play a pivotal role in innate immune responses. TLRs recognize conserved motifs found in various pathogens and mediate defense responses. Triggering of the TLR pathway leads to the activation of NF- κ B and subsequent regulation of immune and inflammatory genes.
Cellular Location	Cytoplasm.
Tissue Location	Ubiquitous.



Western Blot - Anti-MyD88 Rabbit mAb [43Q70G58]

All lanes: R014035 at 1:1,000 dilution

Lane 1: HeL (Human erythroLeukemia suspension cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: A431 (Human epidermoid teratoma cell line) whole cell lysates

Lane 5: T24 (Human bladder cancer epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 33 kDa

Observed band size: 33 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-MyD88 Rabbit mAb [43Q70G58]


Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R014035 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).



Immunohistochemistry - Anti-MyD88 Rabbit mAb [43Q70G58]

Sample: Paraformaldehyde-fixed, paraffin embedded human hepatoma cancer tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R014035 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-MyD88 Rabbit mAb [43Q70G58]

Sample: Paraformaldehyde-fixed, paraffin embedded human colorectal carcinoma tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R014035 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.