

# Anti-Phospho-PKC alpha (Thr497) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R014123

## Product Information

Application	ELISA, WB, IHC-P/IF (Tissue-P)
Reactivity	Rat, Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	37E95G49
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Thr497 of human PKC alpha
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-PKC alpha (Thr497) Rabbit mAb [37E95G49] is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Synonyms	AAG6, Aging associated gene 6, aPKC, KPCA_HUMAN, PKC alpha, PKC-A, PKC-alpha, PKCA, PRKACA, PRKCA, Protein Kinase C alpha, Protein kinase C alpha type.
Calculated MW	Calculated MW: 77 kDa; Observed MW: 77 kDa
Uniprot ID	P17252, P20444, P05696
Gene ID	5578, 18750, 24680
Background	Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This kinase has been reported to play roles in many different cellular processes, such as cell adhesion, cell transformation, cell cycle checkpoint, and cell volume control. Knockout studies in mice suggest that this kinase may be a fundamental regulator of cardiac contractility and Ca(2+) handling in myocytes. [provided by RefSeq, Jul 2008]
Cellular Location	Cytoplasm. Cell membrane. Nucleus.



Western Blot - Anti-Phospho-PKC alpha (Thr497) Rabbit mAb [37E95G49]

All lanes: R014123 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: A431 (Human epidermoid teratoma cell line) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 77 kDa

Observed band size: 77 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-Phospho-PKC alpha (Thr497) Rabbit mAb [37E95G49]

Sample: Paraformaldehyde-fixed, paraffin embedded rat colon tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R014123 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Phospho-PKC alpha (Thr497) Rabbit mAb [37E95G49]

Sample: Paraformaldehyde-fixed, paraffin embedded human endometrial carcinoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R014123 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Phospho-PKC alpha (Thr497) Rabbit mAb [37E95G49]

Sample: Paraformaldehyde-fixed, paraffin embedded mouse liver tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R014123 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.