

Anti-Phospho-STAT1 (Tyr701) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R014159

Product Information

Application	WB, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000
Host	Rabbit
Clonality	Monoclonal
Clone No.	62J91S75
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human STAT1
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-STAT1 (Tyr701) Rabbit mAb [62J91S75] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Signal transducer and activator of transcription 1 91kD, CANDF7, DKFZp686B04100, IMD31A, IMD31B, IMD31C, ISGF 3, ISGF-3, OTTHUMP00000163552, OTTHUMP00000165046, OTTHUMP00000165047, OTTHUMP00000205845, Signal transducer and activator of transcription 1 91kDa, Signal transducer and activator of transcription 1, Signal transducer and activator of transcription 1, 91kD, Signal transducer and activator of transcription 1-alpha/beta, STAT 1, Stat1, STAT1_HUMAN, STAT91, Transcription factor ISGF 3 components p91 p84, Transcription factor ISGF-3 components p91/p84, Transcription factor ISGF3 components p91/p84, XStat1.
Calculated MW	Calculated MW: 87 kDa; Observed MW: 87 kDa
Uniprot ID	P42224
Gene ID	6772
Background	The Stat1 transcription factor is activated in response to a large number of ligands and is essential for responsiveness to IFN- α and IFN- γ . Stat1 protein exists as a pair of isoforms, Stat1 α (91 kDa) and the splice variant Stat1 β (84 kDa). In most cells, both isoforms are activated by IFN- α , but only Stat1 α is activated by IFN- γ . The inappropriate activation of Stat1 occurs in many tumors.
Cellular Location	Cytoplasm. Nucleus. Translocated into the nucleus in response to IFN-gamma-induced tyrosine phosphorylation and dimerization.



Western Blot - Anti-Phospho-STAT1 (Tyr701) Rabbit mAb [62J91S75]

All lanes: R014159 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: Mouse small intestine whole tissue lysates

Lane 3: Rat heart whole tissue lysates

Lane 4: Rat spleen whole tissue lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 87 kDa

Observed band size: 87 kDa

Developed using the ECL technique (Cat. No. SQ201).



Western Blot - Anti-Phospho-STAT1 (Tyr701) Rabbit mAb [62J91S75]

All lanes: R014159 at 1:1,000 dilution

Lane 1: Rat heart whole tissue lysates

Lane 2: Rat spleen whole tissue lysates

Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 87 kDa

Observed band size: 87 kDa

Developed using the ECL technique (Cat. No. SQ201).