

Anti-CD127 Mouse mAb

Purified Recombinant Mouse Monoclonal Antibody

Catalog # M900028


Product Information

Application	ELISA, FC
Reactivity	Human
Dilution	FC 1:100~1:500
Host	Mouse
Clonality	Monoclonal
Clone No.	68C65P27
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human CD127
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-CD127 Mouse mAb [68C65P27] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	CD 127, CD127, CD127 antigen, CDw127, IL 7R alpha, IL 7R, IL-7 receptor subunit alpha, IL-7R subunit alpha, IL-7R-alpha, IL-7RA, IL7R, IL7RA, IL7RA_HUMAN, IL7Ralpha, ILRA, Interleukin 7 receptor alpha chain, Interleukin 7 receptor, Interleukin 7 receptor isoform H5 6, Interleukin-7 receptor subunit alpha.
Uniprot ID	P16871
Gene ID	3575
Background	CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor α chain or IL-7R α . It forms a heterodimer with the common γ chain (γ c or CD132) which is shared with the receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells through early pre-B stage cells, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T cells, and bone marrow stromal cells. CD127 has been reported to be a useful marker for identifying memory and effector T cells. Studies have shown that CD127 expression is down-modulated on Treg cells. It can be used as a marker for differentiation of Treg and conventional T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cell proliferation and development.
Cellular Location	Secreted and Cell membrane.

Validation Images



Human PBMCs were stained with PerCP-Cy5.5-M900028.