

Anti-Phospho-RSK1 (Ser380) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R014295

Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	72J08F97
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Phospho-RSK1(S380)
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-RSK1 (Ser380) Rabbit mAb [72J08F97] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	90 kDa ribosomal protein S6 kinase 1, dJ590P13.1 (ribosomal protein S6 kinase, 90kD, polypeptide 1, dJ590P13.1, EC 2.7.11.1, HU 1, HU1, KS6A1_HUMAN, MAP kinase activated protein kinase 1a, MAP kinase-activated protein kinase 1a, MAPK-activated protein kinase 1a, MAPKAP kinase 1a, MAPKAPK-1a, MAPKAPK1A, MGC79981, Mitogen-activated protein kinase-activated protein kinase 1A, OTTHUMP00000004113, p90 RSK1, p90-RSK 1, p90rsk, p90RSK1, p90S6K, pp90RSK1, Ribosomal protein S6 kinase 90kD 1, Ribosomal protein S6 kinase 90kD polypeptide 1, Ribosomal protein S6 kinase 90kDa polypeptide 1, Ribosomal protein S6 kinase alpha 1, Ribosomal protein S6 kinase alpha-1, Ribosomal protein S6 kinase polypeptide 1, Ribosomal S6 kinase 1, RPS6K1 alpha, rps6ka, Rps6ka1, RSK 1, RSK 1 p90, RSK, RSK-1, RSK1, RSK1p90, S6K alpha 1.
Calculated MW	Calculated MW: 83 kDa; Observed MW: 90 kDa
Uniprot ID	Q15418
Gene ID	6195
Background	Rsk1 is a member of a family of 90kDa ribosomal protein S6 kinases, which includes Rsk1, Rsk2 and Rsk3. These are broadly expressed serine / threonine protein kinases activated in response to mitogenic stimuli, including extracellular signal regulated protein kinases Erk1 and Erk2. Rsk1 is activated by MAPK in vitro and in vivo via phosphorylation. Active Rsk1s appear to play a major role in transcriptional regulation by translocating to the nucleus and phosphorylating c-Fos and CREB.



Western Blot - Anti-Phospho-RSK1 (Ser380) Rabbit mAb [72J08F97]

All lanes: R014295 at 1:1,000 dilution

Lane 1: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 2: C2C12 (Mouse myoblasts epithelial cell) whole cell lysates

Lane 3: A431 (Human epidermoid teratoma cell line) whole cell lysates

Lane 4: T24 (Human bladder cancer epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 83 kDa

Observed band size: 90 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-Phospho-RSK1 (Ser380) Rabbit mAb [72J08F97]

Sample: Paraformaldehyde-fixed, paraffin embedded human gastric cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R014295 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-Phospho-RSK1 (Ser380) Rabbit mAb [72J08F97]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5%

Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R014295 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).