

Anti-ACADVL/VLCAD Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R014300

Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	76G91A35
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human ACADVL/VLCAD
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-ACADVL/VLCAD Rabbit mAb [76G91A35] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	ACAD 6, ACAD6, ACADV_HUMAN, Acadvl, Acyl CoA dehydrogenase very long chain, Acyl Coenzyme A dehydrogenase very long chain, LCACD, mitochondrial, Very long chain specific acyl CoA dehydrogenase, Very long chain specific acyl CoA dehydrogenase mitochondrial, Very long-chain specific acyl-CoA dehydrogenase, VLCAD.
Calculated MW	Calculated MW: 70 kDa; Observed MW: 65 kDa
Uniprot ID	P49748
Gene ID	37
Background	Active toward esters of long-chain and very long chain fatty acids such as palmitoyl-CoA, myristoyl-CoA and stearoyl-CoA. Can accommodate substrate acyl chain lengths as long as 24 carbons, but shows little activity for substrates of less than 12 carbons.
Cellular Location	Mitochondrion inner membrane.



Western Blot - Anti-ACADVL/VLCAD Rabbit mAb [76G91A35]

All lanes: R014300 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: A431 (Human epidermoid teratoma cell line) whole cell lysates

Lane 5: T24 (Human bladder cancer epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 70 kDa

Observed band size: 65 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-ACADVL/VLCAD Rabbit mAb [76G91A35]


Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R014300 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).



Immunohistochemistry - Anti-ACADVL/VLCAD Rabbit mAb [76G91A35]

Sample: Paraformaldehyde-fixed, paraffin embedded human colonic cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R014300 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-ACADVL/VLCAD Rabbit mAb [76G91A35]

Sample: Paraformaldehyde-fixed, paraffin embedded human renal carcinoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R014300 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.