

Anti-ACADVL/VLCAD Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R014770

Product Information

Application	ELISA, WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC
Reactivity	Human
Dilution	WB 1:1,000~1:3,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	91N53M39
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human ACADVL
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-ACADVL/VLCAD Rabbit mAb [91N53M39] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	ACAD 6, ACAD6, ACADV_HUMAN, Acadvl, Acyl CoA dehydrogenase very long chain, Acyl Coenzyme A dehydrogenase very long chain, LCACD, mitochondrial, Very long chain specific acyl CoA dehydrogenase, Very long chain specific acyl CoA dehydrogenase mitochondrial, Very long-chain specific acyl-CoA dehydrogenase, VLCAD.
Calculated MW	Calculated MW: 70 kDa; Observed MW: 70 kDa
Uniprot ID	P49748
Gene ID	37
Background	The protein encoded by this gene is targeted to the inner mitochondrial membrane where it catalyzes the first step of the mitochondrial fatty acid beta-oxidation pathway. This acyl-Coenzyme A dehydrogenase is specific to long-chain and very-long-chain fatty acids. A deficiency in this gene product reduces myocardial fatty acid beta-oxidation and is associated with cardiomyopathy. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]
Cellular Location	Mitochondrion inner membrane.



Western Blot - Anti-ACADVL/VLCAD Rabbit mAb [91N53M39]

All lanes: R014770 at 1:3,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: U2OS (Human osteosarcoma epithelial cell) whole cell lysates

Lane 5: T24 (Human bladder cancer epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 70 kDa

Observed band size: 70 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-ACADVL/VLCAD Rabbit mAb [91N53M39]

Sample: Paraformaldehyde-fixed, paraffin embedded human renal carcinoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R014770 at 1:100 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-ACADVL/VLCAD Rabbit mAb [91N53M39]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R014770 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).