

# Anti-Fatty Acid Synthase Mouse mAb

Purified Recombinant Mouse Monoclonal Antibody


Catalog # M014824

## Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human
Dilution	WB 1:1,000~1:8,000; IHC-P 1:100~1:300; IF 1:100~1:200
Host	Mouse
Clonality	Monoclonal
Clone No.	39M66G65
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Fatty Acid Synthase
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Fatty Acid Synthase Mouse mAb [39M66G65] is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Synonyms	[Acyl-carrier-protein] S acetyltransferase, [Acyl-carrier-protein] S malonyltransferase, 3-hydroxypalmitoyl-[acyl-carrier-protein] dehydratase, 3-oxoacyl-[acyl-carrier-protein] reductase, 3-oxoacyl-[acyl-carrier-protein] synthase, Enoyl-[acyl-carrier-protein] reductase, FAS, FAS_HUMAN, FASN, Fatty acid synthase, MGC14367, MGC15706, OA 519, Oleoyl-[acyl-carrier-protein] hydrolase, SDR27X1, Short chain dehydrogenase/reductase family 27X member 1.
Calculated MW	Calculated MW: 273 kDa; Observed MW: 273 kDa
Uniprot ID	P49327
Gene ID	2194
Background	The enzyme encoded by this gene is a multifunctional protein. Its main function is to catalyze the synthesis of palmitate from acetyl-CoA and malonyl-CoA, in the presence of NADPH, into long-chain saturated fatty acids. In some cancer cell lines, this protein has been found to be fused with estrogen receptor-alpha (ER-alpha), in which the N-terminus of FAS is fused in-frame with the C-terminus of ER-alpha. [provided by RefSeq, Jul 2008]
Cellular Location	Cytoplasm. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.
Tissue Location	Ubiquitous. Prominent expression in brain, lung, and liver.



Western Blot - Anti-Fatty Acid Synthase Mouse mAb [39M66G65]

All lanes: M014824 at 1:8,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: Caco2 (Human colorectal adenocarcinoma epithelial cell) whole cell lysates

Lane 5: 293T (Human embryonic kidney cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Mouse IgG (H+L), HRP Conjugated (Cat. No. LF101) at 1:5,000 dilution

Predicted band size: 273 kDa

Observed band size: 273 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-Fatty Acid Synthase Mouse mAb [39M66G65]


Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibody: M014824 at 1:100 dilution

Secondary antibody: Goat anti-Mouse (488) at 1:1,000 dilution (shown in green)

Nuclei were stained with DAPI (shown in blue).



Immunohistochemistry - Anti-Fatty Acid Synthase Mouse mAb [39M66G65]

Sample: Paraformaldehyde-fixed, paraffin embedded human hepatoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: M014824 at 1:300 dilution

Secondary antibody: Goat Anti-Mouse IgG (H+L), HRP conjugated at 1:1,000 dilution  
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Fatty Acid Synthase Mouse mAb [39M66G65]

Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: M014824 at 1:300 dilution

Secondary antibody: Goat Anti-Mouse IgG (H+L), HRP conjugated at 1:1,000 dilution  
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.