

Anti-p53 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R014950

Product Information

Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	88T68J59
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Mutant p53
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-p53 Rabbit mAb [88T68J59] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Antigen NY-CO-13, BCC7, Cellular tumor antigen p53, FLJ92943, LFS1, Mutant tumor protein 53, p53, p53 tumor suppressor, P53_HUMAN, Phosphoprotein p53, Tp53, Transformation related protein 53, TRP53, tumor antigen p55, Tumor protein 53, Tumor protein p53, Tumor suppressor p53.
Calculated MW	Calculated MW: 44 kDa; Observed MW: 55 kDa
Uniprot ID	P04637
Gene ID	7157
Background	This gene encodes tumor protein p53, which responds to diverse cellular stresses to regulate target genes that induce cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. p53 protein is expressed at low level in normal cells and at a high level in a variety of transformed cell lines, where it's believed to contribute to transformation and malignancy. p53 is a DNA-binding protein containing transcription activation, DNA-binding, and oligomerization domains. It is postulated to bind to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion, and thus function as a tumor suppressor. Mice deficient for this gene are developmentally normal but are susceptible to spontaneous tumors. Evidence to date shows that this gene contains one promoter, in contrast to alternative promoters of the human gene, and transcribes a few of splice variants which encode different isoforms, although the biological validity or the full-length nature of some variants has not been determined. [provided by RefSeq, Jul 2008]
Cellular Location	Cytoplasm; Cytoplasm. Nucleus. Nucleus > PML body. Endoplasmic reticulum. Interaction with BANP promotes nuclear localization. Recruited into PML bodies together with CHEK2; Nucleus. Cytoplasm. Localized in both nucleus and cytoplasm in most cells. In some cells. forms foci in the nucleus that are different from nucleoli: Nucleus. Cytoplasm. Localized in the



Western Blot - Anti-p53 Rabbit mAb [88T68J59]

All lanes: R014950 at 1:1,000 dilution

Lane 1: SW620 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 2: 293T (Human embryonic kidney cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 44 kDa

Observed band size: 55 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-p53 Rabbit mAb [88T68J59]

Sample: Paraformaldehyde-fixed, paraffin embedded human colonic cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R014946 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.