

Anti-Phospho-Chk1 (Ser280) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R015019

Product Information

Application	WB, IF (Cell)/ICC, ELISA
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	30I72T48
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Phospho-Chk1 (S280)
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-Chk1 (Ser280) Rabbit mAb [30I72T48] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	C85740, Cell cycle checkpoint kinase, Checkpoint, <i>S. pombe</i> , homolog of, 1, Checkpoint kinase 1, Checkpoint kinase 1 homolog (<i>S. pombe</i>), CHEK 1, Chk1, Chk 1, Chk1, CHK1 checkpoint homolog (<i>S. pombe</i>), CHK1_HUMAN, EC 2.7.11.1, rad27, Serine/threonine protein kinase Chk1, Serine/threonine-protein kinase CHK1, STT3, subunit of the oligosaccharyltransferase complex, homolog A (<i>S. cerevisiae</i>).
Calculated MW	Calculated MW: 54 kDa; Observed MW: 54 kDa
Uniprot ID	O14757
Gene ID	1111
Background	The protein encoded by this gene belongs to the Ser/Thr protein kinase family. It is required for checkpoint mediated cell cycle arrest in response to DNA damage or the presence of unreplicated DNA. This protein acts to integrate signals from ATM and ATR, two cell cycle proteins involved in DNA damage responses, that also associate with chromatin in meiotic prophase I. Phosphorylation of CDC25A protein phosphatase by this protein is required for cells to delay cell cycle progression in response to double-strand DNA breaks. Several alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Oct 2011]



Western Blot - Anti-Phospho-Chk1 (Ser280) Rabbit mAb [30I72T48]

All lanes: R015019 at 1:1,000 dilution

Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2 : HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3 : HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 54 kDa

Observed band size: 54 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-Phospho-Chk1 (Ser280) Rabbit mAb [30I72T48]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R015019 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).