

Anti-PRAS40 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R015160

Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:2,000~1:10,000
Host	Rabbit
Clonality	Monoclonal
Clone No.	56A30P46
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human PRAS40
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-PRAS40 Rabbit mAb [56A30P46] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	40 kDa proline rich AKT substrate, 40 kDa proline-rich AKT substrate, AKT1 S1, AKT1 substrate 1 (proline rich), AKT1 substrate 1, AKT1S 1, AKT1S1, AKTS1_HUMAN, Lobe, MGC2865, PRAS 40, PRAS, PRAS40, Proline rich akt substrate, Proline rich Akt substrate 40 kDa, Proline-rich AKT1 substrate 1.
Calculated MW	Calculated MW: 27 kDa; Observed MW: 40 kDa
Uniprot ID	Q96B36
Gene ID	84335
Background	AKT1S1 is a proline-rich substrate of AKT (MIM 164730) that binds 14-3-3 protein (see YWHAH, MIM 113508) when phosphorylated (Kovacina et al., 2003 [PubMed 12524439]).[supplied by OMIM, Mar 2008]
Cellular Location	Cytoplasm > cytosol. Found in the cytosolic fraction of the brain.
Tissue Location	Widely expressed with highest levels of expression in liver and heart. Expressed at higher levels in cancer cell lines (e.g. A549 and HeLa) than in normal cell lines (e.g. HEK293).



Western Blot - Anti-PRAS40 Rabbit mAb [56A30P46]

All lanes: R015160 at 1:5,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: SW620 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 5: Caco2 (Human colorectal adenocarcinoma epithelial cell) whole cell lysates

Lane 6: Jurkat (Human T lymphocytic leukemia cell) whole cell lysates

Lane 7: 293T (Human embryonic kidney cell) whole cell lysates

Lane 8: SCC-9 (Human tongue squamous carcinoma epithelial cell) whole cell lysates

Lane 9: C2C12 (Mouse myoblasts epithelial cell) whole cell lysates

Lane 10: Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates

Lane 11: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates

Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at

1:5,000 dilution

Predicted band size: 27 kDa

Observed band size: 40 kDa

Developed using the ECL technique (Cat. No. SQ201).