

PE Mouse Anti-Human CD127

Purified PE-conjugated Recombinant Mouse Monoclonal Antibody


Catalog # F101208

Product Information


Application	FC
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone No.	68C65P27
Isotype	IgG
Label	PE
Immunogen	Recombinant protein of human CD127
Format	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA.
Storage	Shipped on wet ice. Store undiluted between 2°C and 8°C and protected from prolonged exposure to light. Do not freeze. Stable for 24 months from date of receipt.
Precautions	PE Mouse Anti-Human CD127 [68C65P27] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	CD 127, CD127, CD127 antigen, CDw127, IL 7R alpha, IL 7R, IL-7 receptor subunit alpha, IL-7R subunit alpha, IL-7R-alpha, IL-7RA, IL7R, IL7RA, IL7RA_HUMAN, IL7Ralpha, ILRA, Interleukin 7 receptor alpha chain, Interleukin 7 receptor, Interleukin 7 receptor isoform H5 6, Interleukin-7 receptor subunit alpha.
Uniprot ID	P16871
Gene ID	3575
Background	CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor α chain or IL-7R α . It forms a heterodimer with the common γ chain (γ c or CD132) which is shared with the receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells through early pre-B stage cells, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T cells, and bone marrow stromal cells. CD127 has been reported to be a useful marker for identifying memory and effector T cells. Studies have shown that CD127 expression is down-modulated on Treg cells. It can be used as a marker for differentiation of Treg and conventional T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cell proliferation and development.
Cellular Location	Secreted and Cell membrane.



Human peripheral blood lymphocytes were stained with PE Mouse Anti-Human CD127 [68C65P27].



Human peripheral blood lymphocytes were stained with PE Mouse Anti-Human CD127 [68C65P27], Blue histogram. Un-stained control is Pink histogram.