

## Anti-Phospho-RPS6 (Ser240/Ser244) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R015190

### Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	95D07D76
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Ser240/Ser244 of human RPS6
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-RPS6 (Ser240/Ser244) Rabbit mAb [95D07D76] is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information

Synonyms	40S ribosomal protein S6; Air8; NP33; Phosphoprotein NP33; Pp30; Ribosomal protein S6; RP S6; rps6; RS6; RS6_HUMAN; S6; S6 Ribosomal Protein.
Calculated MW	Calculated MW: 29 kDa; Observed MW: 36 kDa
Uniprot ID	P62753
Gene ID	6194
Background	Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 40S subunit. The protein belongs to the S6E family of ribosomal proteins. It is the major substrate of protein kinases in the ribosome, with subsets of five C-terminal serine residues phosphorylated by different protein kinases. Phosphorylation is induced by a wide range of stimuli, including growth factors, tumor-promoting agents, and mitogens. Dephosphorylation occurs at growth arrest. The protein may contribute to the control of cell growth and proliferation through the selective translation of particular classes of mRNA. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Jul 2008]



Western Blot - Anti-Phospho-RPS6 (Ser240/Ser244) Rabbit mAb [95D07D76]

All lanes: R015190 at 1:1,000 dilution


- Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates
  - Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates
  - Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates
  - Lane 4: SW620 (Human colorectal carcinoma epithelial cell) whole cell lysates
  - Lane 5: Caco2 (Human colorectal adenocarcinoma epithelial cell) whole cell lysates
  - Lane 6: Jurkat (Human T lymphocytic leukemia cell) whole cell lysates
  - Lane 7: 293T (Human embryonic kidney cell) whole cell lysates
  - Lane 8: SCC-9 (Human tongue squamous carcinoma epithelial cell) whole cell lysates
  - Lane 9: C2C12 (Mouse myoblasts epithelial cell) whole cell lysates
  - Lane 10: Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates
  - Lane 11: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates
- Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 29 kDa

Observed band size: 36 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-Phospho-RPS6 (Ser240/Ser244) Rabbit mAb [95D07D76]


Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R015190 at 1:100 dilution and  $\alpha$ -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).



Immunohistochemistry - Anti-Phospho-RPS6 (Ser240/Ser244) Rabbit mAb [95D07D76]

Sample: Paraformaldehyde-fixed, paraffin embedded rat brain tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R015190 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution  
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Phospho-RPS6 (Ser240/Ser244) Rabbit mAb [95D07D76]

Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R015190 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution  
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Phospho-RPS6 (Ser240/Ser244) Rabbit mAb [95D07D76]

Sample: Paraformaldehyde-fixed, paraffin embedded mouse stomach tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R015190 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.