

Anti-Epac1 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R015460

Product Information

Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Mouse, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	57T72D78
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Epac 1
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Epac1 Rabbit mAb [57T72D78] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	bcm910; CAMP GEF1; cAMP regulated guanine nucleotide exchange factor I; CAMPGEF1; CGEF 1; CGEF1; EPA1; Epac 1; EPAC; EPAC1; Exchange factor directly activated by cAMP 1; Exchange protein directly activated by cAMP 1; MGC21410; RAP guanine nucleotide exchange factor; Rap guanine nucleotide exchange factor (GEF) 3; RAP guanine nucleotide exchange factor 3; Rap1 guanine nucleotide exchange factor directly activated by cAMP; RAPGEF3.
Calculated MW	Calculated MW: 104 kDa; Observed MW: 104 kDa
Uniprot ID	O95398
Gene ID	10411
Background	Guanine nucleotide exchange factor (GEF) for RAP1A and RAP2A small GTPases that is activated by binding cAMP. Through simultaneous binding of PDE3B to RAPGEF3 and PIK3R6 is assembled in a signaling complex in which it activates the PI3K gamma complex and which is involved in angiogenesis. Plays a role in the modulation of the cAMP-induced dynamic control of endothelial barrier function through a pathway that is independent on Rho-mediated signaling. Required for the actin rearrangement at cell-cell junctions, such as stress fibers and junctional actin.
Cellular Location	Endomembrane system



Western Blot - Anti-Epac1 Rabbit mAb [57T72D78]

All lanes: R015460 at 1:1,000 dilution

Lane 1: Mouse brain whole tissue lysates

Lane 2: Rat brain whole tissue lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 104 kDa

Observed band size: 104 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-Epac1 Rabbit mAb [57T72D78]

Sample: Paraformaldehyde-fixed, paraffin embedded rat kidney tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R015460 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Epac1 Rabbit mAb [57T72D78]

Sample: Paraformaldehyde-fixed, paraffin embedded mouse ovary tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R015460 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.