

Anti-STAT4 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R015470

Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	37J76R51
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human STAT4
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-STAT4 Rabbit mAb [37J76R51] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Signal transducer and activator of transcription 4; SLEB11; STAT4; STAT4_HUMAN.
Calculated MW	Calculated MW: 86 kDa; Observed MW: 86 kDa
Uniprot ID	Q14765
Gene ID	6775
Background	The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is essential for mediating responses to IL12 in lymphocytes, and regulating the differentiation of T helper cells. Mutations in this gene may be associated with systemic lupus erythematosus and rheumatoid arthritis. Alternate splicing results in multiple transcript variants that encode the same protein. [provided by RefSeq, Aug 2011]
Cellular Location	Cytoplasm. Nucleus. Translocated into the nucleus in response to phosphorylation.



Western Blot - Anti-STAT4 Rabbit mAb [37J76R51]

All lanes: R015470 at 1:1,000 dilution

Lane 1: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 86 kDa

Observed band size: 86 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-STAT4 Rabbit mAb [37J76R51]

Sample: Paraformaldehyde-fixed, paraffin embedded human cervical cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R015470 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-STAT4 Rabbit mAb [37J76R51]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R015470 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).