

Recombinant Mouse IL-1 alpha

Catalog # FL071

Product Specifications

| | |
|---------------------|---|
| Appearance | Sterile filtered White lyophilized (freeze-dried) powder. |
| Purity | > 97% by SDS-PAGE or HPLC. |
| Endotoxin | < 0.01 EU/ μ g of rMuIL-1 α protein as determined by LAL method. |
| Expression System | Expressed in E. coli. |
| Species | Mouse |
| Tag | Tag free. |
| Activity | Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using mouse D10S cells is less than 2.0 pg/ml, corresponding to a specific activity of $\geq 5.0 \times 10^8$ IU/mg. |
| Formulation | Lyophilized from a 0.2 μ m filtered concentrated solution in PBS, pH 7.4. |
| Reconstitution | Before use this product, please read the direction below carefully. This vial must be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in a sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at $\leq -20^\circ\text{C}$. Further dilutions should be made in appropriate buffered solutions. |
| Accession # | Q3U0Y6 Ser115-Ser270 |
| Amino acid sequence | SAPYTYQSDLRYKLMKLVKQKFMNDSLNTIYQDVVKHYLSTTWLNDLQQEVKFDMYAYSSGGDDSKYPVTLKISDSQLFVSAQGEDQPVLLKELPETPKLITGSETDLIFFWKSINSKNYFTSAAYPELFIATKEQSRVHLARGLPSMTDFQIS |
| Molecular weight | Approximately 17.9 kDa, a single non-glycosylated polypeptide chain containing 156 amino acids. |
| Synonyms | Lymphocyte-activating factor, LAF, Leukocyte Endogenous Mediator, LEM, Endogenous Pyrogen, EP, Mononuclear Cell Factor, MCF, Hematopoietin-1 |
| Stability & Storage | Shipped on wet ice. For long term storage, the product should be stored $\leq -20^\circ\text{C}$. Please avoid repeated freeze-thaw cycles after reconstitution. 36 months from date of receipt, -20 to -70°C as supplied. 1 month, 2 to 8°C under sterile conditions after reconstitution. 3 months, -20 to -70°C under sterile conditions after reconstitution. |
| Precautions | Recombinant Mouse IL-1 alpha is for research use only and not for use in diagnostic or therapeutic procedures. |

Background

IL-1 α is a non-secreted, proinflammatory cytokine produced in a variety of cells, including monocytes, tissue macrophages, keratinocytes, and other epithelial cells. It possesses metabolic, physiological, hematopoietic activities, and plays one of the central roles in the regulation of the immune responses. Both IL-1 α and IL-1 β bind to the same receptor and have similar, if not identical, biological properties. These cytokines have a broad range of activities including the stimulation of thymocyte proliferation by inducing IL-2 release, B cell maturation and proliferation, mitogenic FGF-like activity and the release of prostaglandin and collagenase from synovial cells. However, whereas IL-1 β is a secreted cytokine, IL-1 α is predominantly a cell-associated cytokine. Recombinant Mouse IL-1 α is a 17.9kDa protein containing 156 amino acid.

