

## Anti-MTTP Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P103875


### Product Information

Application	WB, IF (Cell)/ICC, ELISA
Reactivity	Mouse, Rat
Dilution	WB 1:500~1:2,000; IF 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 550-650 of human MTTP (NP_000244.2).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-MTTP Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.


### Protein Information

Synonyms	ABL; MTP; MTTP.
Calculated MW	Calculated MW: 99 kDa; Observed MW: 99 kDa
Uniprot ID	P55157
Gene ID	4547
Background	MTP encodes the large subunit of the heterodimeric microsomal triglyceride transfer protein. Protein disulfide isomerase (PDI) completes the heterodimeric microsomal triglyceride transfer protein, which has been shown to play a central role in lipoprotein assembly. Mutations in MTP can cause abetalipoproteinemia.

## Validation Images



Immunofluorescence analysis of NIH/3T3 cells using MTTP Rabbit pAb (P103875) at dilution of 1:20 (40 $\times$  lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using MTTP Rabbit pAb (P103875) at 1:1,000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25 $\mu$ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit.

Exposure time: 90s.