

Anti-ULK1/ULK2 Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P104389


Product Information

Application	WB, ELISA
Reactivity	Human
Dilution	WB 1:500~1:2,000
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-283 of human ULK1/ULK2 (NP_003556.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-ULK1/ULK2 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	ATG 1; ATG1; ATG1 autophagy related 1 homolog; ATG1A; Autophagy related protein 1 homolog; Autophagy-related protein 1 homolog; FLJ38455; FLJ46475; hATG1; KIAA0722; Serine/threonine protein kinase ULK1; Serine/threonine protein kinase Unc51.1; Serine/threonine-protein kinase ULK1; ULK 1; ULK1; ULK1_HUMAN; Unc 51 (C. elegans) like kinase 1; UNC 51; Unc 51 like kinase 1; Unc-51 like kinase 1 (C. elegans) ; Unc-51-like kinase 1; UNC51; UNC51, C. elegans, homolog of; Unc51.1; ATG1B; KIAA0623; Serine/threonine protein kinase ULK2; Serine/threonine-protein kinase ULK2; ULK2; ULK2_HUMAN; Unc 51 (C. elegans) like kinase 2; Unc 51 like autophagy activating kinase 2; Unc 51 like kinase 2; Unc-51-like kinase 2; Unc51.2.
Calculated MW	Observed MW: 113 kDa
Uniprot ID	O75385, Q8IYT8
Gene ID	8408, 9706
Background	Mammalian ULK1 (unc-51 like kinase 1) and ULK2, <i>Caenorhabditis elegans</i> UNC-51, and <i>Drosophila melanogaster</i> Atg1 are serine/threonine kinases that regulate flux through the autophagy pathway in response to various types of cellular stress. ULK1 and ULK2 all have different tissue-specific expressions partially contributing to diverse and ULK-specific interaction networks in various tissues. ULK1 (or its close family member ULK2) interacts with several other proteins to form a large complex that is required for the formation of an autophagosome. The other proteins that complex with ULK1/ULK2 include ATG13, ATG101, and RB1CC1 (FIP200/ATG17). AMPK, activated during low nutrient conditions, directly phosphorylates ULK1 at multiple sites including Ser317, Ser555, and Ser777. Conversely, mTOR, which is a regulator of cell growth and is an inhibitor of autophagy, phosphorylates ULK1 at Ser757 and disrupts the interaction between ULK1 and AMPK.

Validation Images



Western blot analysis of various lysates using ULK1/ULK2 Rabbit pAb (P104389) at 1:500 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit.

Exposure time: 90s.