

Anti-IL18 Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P104517

Product Information


Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:1,000; IHC-P 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 36-192 of mouse IL18 (NP_032386.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-IL18 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Synonyms	Ig1f; IL-18; IL18.
Calculated MW	Calculated MW: 22 kDa; Observed MW: 22 kDa/18 kDa
Uniprot ID	P70380
Gene ID	16173
Background	Enables cytokine activity. Involved in lipid homeostasis; positive regulation of cell differentiation; and positive regulation of cold-induced thermogenesis. Acts upstream of or within several processes, including interleukin-18-mediated signaling pathway; positive regulation of NIK/NF-kappaB signaling; and positive regulation of interferon-gamma production. Located in extracellular space. Is expressed in several structures, including back skin; dorsal aorta; gut; lymphatic system; and reproductive system. Used to study atopic dermatitis. Human ortholog(s) of this gene implicated in several diseases, including Kawasaki disease; allergic rhinitis; autoimmune disease (multiple); biliary atresia; and hepatitis B. Orthologous to human IL18 (interleukin 18).




Immunohistochemistry analysis of paraffin-embedded Human tonsil using IL18 Rabbit pAb (P104517) at dilution of 1:50 (40 \times lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse spleen using IL18 Rabbit pAb (P104517) at dilution of 1:50 (40 \times lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat spleen using IL18 Rabbit pAb (P104517) at dilution of 1:50 (40 \times lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Western blot analysis of various lysates using IL18 Rabbit pAb (P104517) at 1:1,000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25 μ g per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Kit.
Exposure time: 30s.



Western blot analysis of lysates from Rat brain, using IL18 Rabbit pAb (P104517) at 1:1,000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit.

Exposure time: 180s.



Western blot analysis of Recombinant Human IL-18 Protein using IL18 Rabbit pAb (P104517) at 1:1,000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 10ng/5ng per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit.

Exposure time: 180s.