

Anti-IL1RN Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P104965


Product Information

| | |
|-------------|---|
| Application | WB, IF (Cell)/ICC, ELISA |
| Reactivity | Human, Mouse |
| Dilution | WB 1:100~1:500; IF 1:50~1:200 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Label | Unconjugated |
| Immunogen | Recombinant fusion protein containing a sequence corresponding to amino acids 27-178 of mouse IL1RN (NP_001034790.1). |
| Format | Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3. |
| Storage | Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage. |
| Precautions | Anti-IL1RN Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures. |


Protein Information

| | |
|---------------|---|
| Synonyms | IL-1ra; F630041P17Rik; IL1RN. |
| Calculated MW | Calculated MW: 20 kDa; Observed MW: 18 kDa |
| Uniprot ID | P25085 |
| Gene ID | 16181 |
| Background | Predicted to enable cytokine receptor binding activity. Acts upstream of or within insulin secretion and lipid metabolic process. Located in extracellular region and vesicle. Human ortholog(s) of this gene implicated in several diseases, including autoimmune disease (multiple); eye disease (multiple); kidney failure (multiple); lung disease (multiple); and vascular disease (multiple). Orthologous to human IL1RN (interleukin 1 receptor antagonist). |

Validation Images



Immunofluorescence analysis of NIH/3T3 cells using IL1RN Rabbit pAb (P104965) at dilution of 1:50 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using IL1RN Rabbit pAb (P104965) at 1:500 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Enhanced Kit.
Exposure time: 180s.