

Anti-DVL3 Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P105857

Product Information


| | |
|-------------|---|
| Application | WB, IHC-P/IF (Tissue-P), ELISA |
| Reactivity | Human, Mouse, Rat |
| Dilution | WB 1:500~1:1,000; IHC-P 1:50~1:200 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Label | Unconjugated |
| Immunogen | Recombinant fusion protein containing a sequence corresponding to amino acids 580-650 of human DVL3 (NP_004414.3). |
| Format | Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3. |
| Storage | Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage. |
| Precautions | Anti-DVL3 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information


| | |
|---------------|--|
| Synonyms | DRS3; DVL3. |
| Calculated MW | Calculated MW: 78 kDa; Observed MW: 93 kDa |
| Uniprot ID | Q92997 |
| Gene ID | 1857 |
| Background | This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. |




Immunohistochemistry analysis of paraffin-embedded Mouse brain using DVL3 Rabbit pAb (P105857) at dilution of 1:100 (40× lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse kidney using DVL3 Rabbit pAb (P105857) at dilution of 1:100 (40× lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain using DVL3 Rabbit pAb (P105857) at dilution of 1:100 (40× lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat ovary using DVL3 Rabbit pAb (P105857) at dilution of 1:100 (40× lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Western blot analysis of various lysates using DVL3 Rabbit pAb (P105857) at 1:1,000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit (SQ201).

Exposure time: 10s.