

# Anti-PP2A Catalytic Beta Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P105871

## Product Information


Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:2,000; IHC-P 1:100~1:200; IF 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-309 of human PP2A Catalytic $\beta$ (NP_001009552.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-PP2A Catalytic Beta Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information


Synonyms	PP2CB; PP2A beta; PP2A Catalytic beta.
Calculated MW	Calculated MW: 36 kDa; Observed MW: 36 kDa
Uniprot ID	P62714
Gene ID	5516
Background	This gene encodes the phosphatase 2A catalytic subunit. Protein phosphatase 2A is one of the four major Ser/Thr phosphatases, and it is implicated in the negative control of cell growth and division. It consists of a common heteromeric core enzyme, which is composed of a catalytic subunit and a constant regulatory subunit, that associates with a variety of regulatory subunits. This gene encodes a beta isoform of the catalytic subunit.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma using PP2A Catalytic  $\beta$  Rabbit pAb (P105871) at dilution of 1:100 (40 $\times$  lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat kidney using PP2A Catalytic  $\beta$  Rabbit pAb (P105871) at dilution of 1:100 (40 $\times$  lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse kidney using PP2A Catalytic  $\beta$  Rabbit pAb (P105871) at dilution of 1:100 (40 $\times$  lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of U-2 OS cells using PP2A Catalytic  $\beta$  Rabbit pAb (P105871) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using PP2A Catalytic  $\beta$  Rabbit pAb (P105871) at 1:1,000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25 $\mu$ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit (SQ201).

Exposure time: 30s.