Anti-BCAP29 Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P106790

Product Information

|	
Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:2,000; IHC-P 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 125-241 of human BCAP29 (NP_061332.2).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-BCAP29 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

|	
Synonyms	B29; BAP29; BCAP29.
Calculated MW	Calculated MW: 28 kDa; Observed MW: 28 kDa
Uniprot ID	Q9UHQ4
Gene ID	55973
Background	Involved in osteoblast differentiation. Located in membrane.

Immunohistochemistry analysis of paraffin-embedded Rat kidney using BCAP29 Rabbit pAb (P106790) at dilution of 1:100 (40× lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Human stomach using BCAP29 Rabbit pAb (P106790) at dilution of 1:100 (40× lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Mouse spleen using BCAP29 Rabbit pAb (P106790) at dilution of 1:100 (40× lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

Western blot analysis of various lysates using BCAP29 Rabbit pAb (P106790) at 1:1,000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25μg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Enhanced Kit (SQ201).
Exposure time: 30s.