

## Anti-CDKN2D Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P106898


### Product Information

Application	WB, IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse
Dilution	WB 1:500~1:1,000; IF 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-166 of human CDKN2D (NP_001791.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-CDKN2D Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.


### Protein Information

Synonyms	p19; INK4D; p19-INK4D; CDKN2D.
Calculated MW	Calculated MW: 18 kDa; Observed MW: 18 kDa
Uniprot ID	P55273
Gene ID	1032
Background	The protein encoded by this gene is a member of the INK4 family of cyclin-dependent kinase inhibitors. This protein has been shown to form a stable complex with CDK4 or CDK6, and prevent the activation of the CDK kinases, thus function as a cell growth regulator that controls cell cycle G1 progression. The abundance of the transcript of this gene was found to oscillate in a cell-cycle dependent manner with the lowest expression at mid G1 and a maximal expression during S phase. The negative regulation of the cell cycle involved in this protein was shown to participate in repressing neuronal proliferation, as well as spermatogenesis. Two alternatively spliced variants of this gene, which encode an identical protein, have been reported.

## Validation Images



Immunofluorescence analysis of NIH-3T3 cells using CDKN2D Rabbit pAb (P106898) at dilution of 1:100 (40 $\times$  lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using CDKN2D Rabbit pAb (P106898) at 1:1,000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates / proteins: 25  $\mu$ g per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Kit (SQ201).

Exposure time: 90s.