

## Anti-IL18BP Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P107021

### Product Information


|             |   |
|-------------|---|
| Application | WB, IF (Cell)/ICC, ELISA  |
| Reactivity  | Human, Mouse  |
| Dilution    | WB 1:1,000~1:5,000; IF 1:50~1:100   |
| Host        | Rabbit  |
| Clonality   | Polyclonal  |
| Isotype     | IgG   |
| Label       | Unconjugated  |
| Immunogen   | Recombinant fusion protein containing a sequence corresponding to amino acids 31-194 of human IL18BP (NP_001138529.1).      |
| Format      | Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.                     |
| Storage     | Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage. |
| Precautions | Anti-IL18BP Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.                    |

### Protein Information


|               |  |
|---------------|--|
| Synonyms      | FVH; IL18BPa; IL18BP.  |
| Calculated MW | Calculated MW: 21 kDa; Observed MW: 44 kDa   |
| Uniprot ID    | O95998   |
| Gene ID       | 10068  |
| Background    | The protein encoded by this gene functions as an inhibitor of the proinflammatory cytokine, IL18. It binds IL18, prevents the binding of IL18 to its receptor, and thus inhibits IL18-induced IFN-gamma production, resulting in reduced T-helper type 1 immune responses. This protein is constitutively expressed and secreted in mononuclear cells. Elevated level of this protein is detected in the intestinal tissues of patients with Crohn's disease. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. |

## Validation Images

---



Immunofluorescence analysis of A549 cells using IL18BP Rabbit pAb (P107021).  
Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of lysates from Mouse thymus, using IL18BP Rabbit pAb (P107021) at 1:2,000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.  
Lysates/proteins: 25 $\mu$ g per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Kit (SQ201).  
Exposure time: 180s.